

Grading

1

Your PRINTED name is: _____

2

3

4

Please circle your recitation:

- 1) T 10 2-131 K. Meszaros 2-333 3-7826 karola
- 2) T 10 2-132 A. Barakat 2-172 3-4470 barakat
- 3) T 11 2-132 A. Barakat 2-172 3-4470 barakat
- 4) T 11 2-131 A. Osorno 2-229 3-1589 aosorno
- 5) T 12 2-132 A. Edelman 2-343 3-7770 edelman
- 6) T 12 2-131 K. Meszaros 2-333 3-7826 karola
- 7) T 1 2-132 A. Edelman 2-343 3-7770 edelman
- 8) T 2 2-132 J. Burns 2-333 3-7826 burns
- 9) T 3 2-132 A. Osorno 2-229 3-1589 aosorno

- 1 (24 pts.) Suppose q_1, q_2, q_3 are orthonormal vectors in \mathbb{R}^3 . Find **all possible values** for these 3 by 3 determinants and explain your thinking in 1 sentence each.

(a) $\det \begin{bmatrix} q_1 & q_2 & q_3 \end{bmatrix} =$

(b) $\det \begin{bmatrix} q_1 + q_2 & q_2 + q_3 & q_3 + q_1 \end{bmatrix} =$

(c) $\det \begin{bmatrix} q_1 & q_2 & q_3 \end{bmatrix}$ times $\det \begin{bmatrix} q_2 & q_3 & q_1 \end{bmatrix} =$

This page intentionally blank.

2 (24 pts.) Suppose we take measurements at the 21 equally spaced times $t = -10, -9, \dots, 9, 10$. All measurements are $b_i = 0$ except that $b_{11} = 1$ at the middle time $t = 0$.

(a) Using least squares, what are the best \hat{C} and \hat{D} to fit those 21 points by a straight line $C + Dt$?

(b) You are projecting the vector b onto what subspace? (*Give a basis.*)
Find a nonzero vector perpendicular to that subspace.

This page intentionally blank.

3 (9 + 12 + 9 pts.) The Gram-Schmidt method produces orthonormal vectors q_1, q_2, q_3 from independent vectors a_1, a_2, a_3 in \mathbb{R}^5 . Put those vectors into the columns of 5 by 3 matrices Q and A .

(a) Give formulas using Q and A for the projection matrices P_Q and P_A onto the column spaces of Q and A .

(b) *Is $P_Q = P_A$ and why? What is P_Q times Q ? What is $\det P_Q$?*

(c) Suppose a_4 is a new vector and a_1, a_2, a_3, a_4 are independent. Which of these (if any) is the new Gram-Schmidt vector q_4 ? (P_A and P_Q from above)

$$\begin{array}{lll}
 \mathbf{1.} & \frac{P_Q a_4}{\|P_Q a_4\|} & \mathbf{2.} \frac{a_4 - \frac{a_4^T a_1}{a_1^T a_1} a_1 - \frac{a_4^T a_2}{a_2^T a_2} a_2 - \frac{a_4^T a_3}{a_3^T a_3} a_3}{\| \text{norm of that vector} \|} & \mathbf{3.} \frac{a_4 - P_A a_4}{\|a_4 - P_A a_4\|}
 \end{array}$$

This page intentionally blank.

- 4 (22 pts.) Suppose a 4 by 4 matrix has the same entry \times throughout its first row and column. The other 9 numbers could be anything like 1, 5, 7, 2, 3, 99, π , e , 4.

$$A = \begin{bmatrix} \times & \times & \times & \times \\ \times & \text{any numbers} & & \\ \times & \text{any numbers} & & \\ \times & \text{any numbers} & & \end{bmatrix}$$

- (a) The determinant of A is a polynomial in \times . What is the largest possible degree of that polynomial? **Explain your answer.**
- (b) If those 9 numbers give the identity matrix I , what is $\det A$? Which values of \times give $\det A = 0$?

$$A = \begin{bmatrix} \times & \times & \times & \times \\ \times & 1 & 0 & 0 \\ \times & 0 & 1 & 0 \\ \times & 0 & 0 & 1 \end{bmatrix}$$

This page intentionally blank.