

18.06 Problem Set 2 - Solutions

Due Wednesday, Feb. 21, 2007 at **4:00 p.m.** in 2-106

Each problem is worth 10 points. The date next to the problem number indicates the lecture in which the material is covered.

Problem 1 Monday 2/12

Use Gauss-Jordan elimination to find the inverse of $A = \begin{bmatrix} 1 & 4 & 2 \\ 1 & 4 & 1 \\ 2 & 10 & 5 \end{bmatrix}$.

Solution 1

$$\begin{aligned} & \left[\begin{array}{ccc|ccc} 1 & 4 & 2 & 1 & 0 & 0 \\ 1 & 4 & 1 & 0 & 1 & 0 \\ 2 & 10 & 5 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 4 & 2 & 1 & 0 & 0 \\ 0 & 0 & -1 & -1 & 1 & 0 \\ 2 & 10 & 5 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 4 & 2 & 1 & 0 & 0 \\ 0 & 0 & -1 & -1 & 1 & 0 \\ 0 & 2 & 1 & -2 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 4 & 2 & 1 & 0 & 0 \\ 0 & 2 & 1 & -2 & 0 & 1 \\ 0 & 0 & -1 & -1 & 1 & 0 \end{array} \right] \\ \rightarrow & \left[\begin{array}{ccc|ccc} 1 & 4 & 2 & 1 & 0 & 0 \\ 0 & 2 & 0 & -3 & 1 & 1 \\ 0 & 0 & -1 & -1 & 1 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 4 & 0 & -1 & 2 & 0 \\ 0 & 2 & 0 & -3 & 1 & 1 \\ 0 & 0 & -1 & -1 & 1 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 5 & 0 & -2 \\ 0 & 2 & 0 & -3 & 1 & 1 \\ 0 & 0 & -1 & -1 & 1 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 5 & 0 & -2 \\ 0 & 1 & 0 & -3/2 & 1/2 & 1/2 \\ 0 & 0 & 1 & 1 & -1 & 0 \end{array} \right] \end{aligned}$$

Therefore, A^{-1} is $\begin{bmatrix} 5 & 0 & -2 \\ -3/2 & 1/2 & 1/2 \\ 1 & -1 & 0 \end{bmatrix}$

Problem 2 Monday 2/12

Do problem 29 of section 2.5 in your book.

Solution 2

- (a) True. This matrix has three pivots maximum.
- (b) False. $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ is not invertible.
- (c) True. A^{-1} has inverse A.
- (d) True. A^2 has inverse $A^{-1} * A^{-1}$.

Problem 3 Monday 2/12

Do problem 32 of section 2.5 in your book.

Solution 3

$$\left(\begin{array}{cccc|cccc} 1 & -1 & 1 & -1 & 1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right)$$

→ add the second row to the first row →

$$\left(\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right)$$

→ add the third row to the second row →

$$\left(\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right)$$

→ add the fourth row to the third row →

$$\left(\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right)$$

Therefore, $A^{-1} = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}$.

For 5 by 5 alternating matrix, the inverse is $\begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$.

Problem 4 Wednesday 2/14

Do problem 11 of section 2.6 in your book.

Solution 4

A is already an upper triangular matrix, so for $A = LU$ decomposition, $L = I$, $U = A = \begin{bmatrix} 2 & 4 & 8 \\ 0 & 3 & 9 \\ 0 & 0 & 7 \end{bmatrix}$.

For $A = LDU$ decomposition, $L = I$, $D = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 7 \end{bmatrix}$, $U = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & 3 \\ 0 & 0 & 1 \end{bmatrix}$.

Problem 5 Wednesday 2/14

Compute the LU decomposition for the matrix

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix}$$

Solution 5

$$\left(\begin{array}{cccc|cccc} 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 2 & 2 & 2 & 0 & 1 & 0 & 0 \\ 1 & 2 & 3 & 3 & 0 & 0 & 1 & 0 \\ 1 & 2 & 3 & 4 & 0 & 0 & 0 & 1 \end{array} \right)$$

→ subtract the first row from the second, third, and fourth row →

$$\left(\begin{array}{cccc|cccc} 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & -1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 2 & -1 & 0 & 1 & 0 \\ 0 & 1 & 2 & 3 & -1 & 0 & 0 & 1 \end{array} \right)$$

→ subtract the second row from the third and fourth row →

$$\left(\begin{array}{cccc|cccc} 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & 2 & 0 & -1 & 0 & 1 \end{array} \right)$$

→ subtract the third row from fourth row →

$$\left(\begin{array}{cccc|cccc} 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & -1 & 1 \end{array} \right)$$

Therefore, $NA = U$ for

$$N = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{bmatrix}$$

$$U = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$A = N^{-1}U$, so

$$L = N^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$

$$U = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

gives the LU decomposition of A .

Problem 6 Wednesday 2/14

Do problem 16 of section 2.6 in your book.

Solution 6

Let $c = \begin{bmatrix} p \\ q \\ r \end{bmatrix}$.

$$Lc = b \leftrightarrow \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} p \\ q \\ r \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \leftrightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} p \\ q \\ r \end{bmatrix} = \begin{bmatrix} 4 \\ 1 \\ 2 \end{bmatrix} \leftrightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} p \\ q \\ r \end{bmatrix} = \begin{bmatrix} 4 \\ 1 \\ 1 \end{bmatrix}$$

Therefore, $c = \begin{bmatrix} 4 \\ 1 \\ 1 \end{bmatrix}$

Let $x = \begin{bmatrix} y \\ z \\ w \end{bmatrix}$.

$$Lc = b \leftrightarrow \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} y \\ z \\ w \end{bmatrix} = \begin{bmatrix} 4 \\ 1 \\ 1 \end{bmatrix} \leftrightarrow \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} y \\ z \\ w \end{bmatrix} = \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \leftrightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} y \\ z \\ w \end{bmatrix} = \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}$$

Therefore, $x = \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}$

Here, $Ax = LUx = Lc = b$ and $A = LU = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 2 \\ 1 & 2 & 3 \end{bmatrix}$.

Problem 7 Friday 2/16

Let $P = P_{45}P_{16}P_{56}P_{12}P_{34}$, where the P_{ij} are permutation matrices of order 6.

Solution 7

(a) What is $P \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \end{bmatrix}$?

$$\begin{aligned} P \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \end{bmatrix} &= P_{45}P_{16}P_{56}P_{12}P_{34} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \end{bmatrix} = P_{45}P_{16}P_{56}P_{12} \begin{bmatrix} 1 \\ 2 \\ 4 \\ 3 \\ 5 \\ 6 \end{bmatrix} = P_{45}P_{16}P_{56} \begin{bmatrix} 1 \\ 2 \\ 4 \\ 3 \\ 5 \\ 6 \end{bmatrix} \\ &= P_{45}P_{16} \begin{bmatrix} 2 \\ 1 \\ 4 \\ 3 \\ 6 \\ 5 \end{bmatrix} = P_{45} \begin{bmatrix} 5 \\ 1 \\ 4 \\ 3 \\ 6 \\ 2 \end{bmatrix} = \begin{bmatrix} 5 \\ 1 \\ 4 \\ 6 \\ 3 \\ 2 \end{bmatrix} \end{aligned}$$

(b) What is P ?

P is the permutation matrix sending

row 1 \rightarrow row 2

row 2 \rightarrow row 6

row 3 \rightarrow row 5

row 4 \rightarrow row 3

row 5 \rightarrow row 1

row 6 \rightarrow row 4

when multiplied on the left.

Therefore $P = \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \end{bmatrix}$

Problem 8 Friday 2/16

Do problem 16 of section 2.7 in your book.

Solution 8

(a) $[A^2 + B^2]$ is symmetric

$$\leftrightarrow [(A^2 + B^2)^T = A^2 + B^2]$$

$$\leftrightarrow [(A^2)^T + (B^2)^T = A^2 + B^2]$$

$$\leftrightarrow [(A^T)^2 + (B^T)^2 = A^2 + B^2].$$

This is true since $A = A^T$ and $B = B^T$.

(b) $[(A + B)(A - B)]$ is symmetric

$$\leftrightarrow \{[(A + B)(A - B)]^T = (A + B)(A - B)\}$$

$$\leftrightarrow [(A - B)^T(A + B)^T = (A + B)(A - B)]$$

$$\leftrightarrow [(A^T - B^T)(A^T + B^T) = (A + B)(A - B)]$$

$$\leftrightarrow [(A - B)(A + B) = (A + B)(A - B)]$$

$$\leftrightarrow [A^2 - BA + AB - B^2 = A^2 + BA - AB - B^2]$$

$$\leftrightarrow [AB = BA]$$

This is not true since A and B do not generally commute.

(c) [ABA is symmetric]

$$\leftrightarrow [(ABA)^T = ABA]$$

$$\leftrightarrow [A^T B^T A^T = ABA]$$

This is true since $A = A^T$ and $B = B^T$.

(c) [ABAB is symmetric]

$$\leftrightarrow [(ABAB)^T = ABAB]$$

$$\leftrightarrow [B^T A^T B^T A^T = ABAB]$$

$$\leftrightarrow [BABA = ABAB]$$

This is not true since A and B do not generally commute.

Problem 9 Friday 2/16

$$\text{Let } A = \begin{bmatrix} 2 & 6 & 5 \\ 3 & 9 & 1 \\ 1 & 2 & 3 \end{bmatrix}.$$

Compute an LU factorization of A if one exists, or find a permutation matrix P such that PA = LU otherwise.

Solution 9

$$\text{For } P = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, PA = \begin{bmatrix} 2 & 6 & 5 \\ 1 & 2 & 3 \\ 3 & 9 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 6 & 5 & 1 & 0 & 0 \\ 1 & 2 & 3 & 0 & 1 & 0 \\ 3 & 9 & 1 & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 6 & 5 & 1 & 0 & 0 \\ 0 & -1 & 1/2 & -1/2 & 1 & 0 \\ 0 & 0 & -13/2 & -3/2 & 0 & 1 \end{bmatrix}$$

$$\text{Therefore, for } N = \begin{bmatrix} 1 & 0 & 0 \\ -1/2 & 1 & 0 \\ -3/2 & 0 & 1 \end{bmatrix}, NPA = \begin{bmatrix} 2 & 6 & 5 \\ 0 & -1 & 1/2 \\ 0 & 0 & -13/2 \end{bmatrix} = U.$$

$$\text{Therefore, } PA = N^{-1}U = \begin{bmatrix} 1 & 0 & 0 \\ 1/2 & 1 & 0 \\ 3/2 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 6 & 5 \\ 0 & -1 & 1/2 \\ 0 & 0 & -13/2 \end{bmatrix} \text{ is the LU decomposition of PA.}$$

Problem 10 Friday 2/16

Are the following statements true or false? For each statement explain why it is true or give a counterexample if it is false. All matrices are assumed to be square.

Solution 10

(a) The product of two upper triangular matrices is an upper triangular matrix.

True.

Suppose that A and B is $n \times n$ upper triangular. Then we have $A_{ij} = 0, B_{ij} = 0$ for $i > j$.

$$AB_{ij} = \sum_{k=1}^n A_{ik} B_{kj} = \sum_{k=1}^i A_{ik} B_{kj} + \sum_{k=i+1}^n A_{ik} B_{kj}$$

For $i > j$, the first term is 0 since $A_{ik} = 0$ for $i > k$, and the second term is 0 since $B_{kj} = 0$ for $k > i > j$. Therefore $AB_{ij} = 0$ for $i > j$, which means AB is upper triangular matrix.

(b) The product of two symmetric matrices is a symmetric matrix.

False.

Even if $A^T = A$, and $B^T = B$ $(AB)^T = B^T A^T = BA$ is not guaranteed to be equal to AB.

(c) The product of two permutation matrices is a permutation matrix.

True.

The composition of two permutations is also a permutation.

(d) The inverse of a lower triangular matrix is lower triangular, if it exists .

True.

If a lower triangular matrix L has an inverse, then diagonal entries are all nonzero. Therefore, in the Gauss-Jordan elimination, we can take the k -th row as a k -th pivot to get rid of entries in k -th column. Since $1, 2, 3, \dots, (k-1)$ th row has entry 0 in k -th column, we only get lower triangular matrices as elimination matrices. From the same reasoning in (a), the product of lower triangular matrices is lower triangular. Therefore, we get lower triangular inverse.