

**Grading**

Your PRINTED name is: \_\_\_\_\_

**1****2****3**

Please circle your recitation: \_\_\_\_\_

r01	T 11	4-159	Ailsa Keating	ailsa
r02	T 11	36-153	Rune Haugseng	haugseng
r03	T 12	4-159	Jennifer Park	jmypark
r04	T 12	36-153	Rune Haugseng	haugseng
r05	T 1	4-153	Dimiter Ostrev	ostrev
r06	T 1	4-159	Uhi Rinn Suh	ursuh
r07	T 1	66-144	Ailsa Keating	ailsa
r08	T 2	66-144	Niels Martin Moller	moller
r09	T 2	4-153	Dimiter Ostrev	ostrev
r10	ESG		Gabrielle Stoy	gstoy

**1 (33 pts.)**

Suppose an  $n \times n$  matrix  $A$  has  $n$  independent eigenvectors  $x_1, \dots, x_n$ . Then you could write the solution to  $\frac{du}{dt} = Au$  in three ways:

$$u(t) = e^{At}u(0), \quad \text{or}$$

$$u(t) = Se^{\Lambda t}S^{-1}u(0), \quad \text{or}$$

$$u(t) = c_1e^{\lambda_1 t}x_1 + \dots + c_n e^{\lambda_n t}x_n.$$

Here,  $S = [x_1 \mid x_2 \mid \dots \mid x_n]$ .

(a) From the definition of the exponential of a matrix, show why  $e^{At}$  is the same as  $Se^{\Lambda t}S^{-1}$ .

(b) How do you find  $c_1, \dots, c_n$  from  $u(0)$  and  $S$ ?

- (c) For this specific equation, write  $u(t)$  in any one of the three forms, using *numbers* not symbols: You can choose which form.

$$\frac{du}{dt} = \begin{bmatrix} 1 & 2 \\ -1 & 4 \end{bmatrix} u, \quad \text{starting from } u(0) = \begin{bmatrix} 4 \\ 3 \end{bmatrix}.$$

*This page intentionally blank.*

**2 (30 pts.)**

This question is about the real matrix

$$A = \begin{bmatrix} 1 & c \\ 1 & -1 \end{bmatrix}, \quad \text{for } c \in \mathbb{R}.$$

- (a) - Find the eigenvalues of  $A$ , depending on  $c$ .  
- For which values of  $c$  does  $A$  have real eigenvalues?

(b) - For one particular value of  $c$ , convince me that  $A$  is similar to both the matrix

$$B = \begin{bmatrix} 2 & 0 \\ 0 & -2 \end{bmatrix},$$

and to the matrix

$$C = \begin{bmatrix} 2 & 2 \\ 0 & -2 \end{bmatrix}.$$

- Don't forget to say which value  $c$  this happens for.

- (c) For one particular value of  $c$ , convince me that  $A$  cannot be diagonalized. It is not similar to a diagonal matrix  $\Lambda$ , when  $c$  has that value.
- Which value  $c$ ?
  - Why not?

*This page intentionally blank.*



**3 (37 pts.)**

(a) Suppose  $A$  is an  $n \times n$  symmetric matrix with eigenvalues  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$ .

- What is the largest number real number  $c$  that can be subtracted from the diagonal entries of  $A$ , so that  $A - cI$  is positive semidefinite?
- Why?

(b) Suppose  $B$  is a matrix with independent columns.

- What is the nullspace  $N(B)$ ?

- Show that  $A = B^T B$  is positive definite. Start by saying what that means about  $x^T A x$ .

(c) This matrix  $A$  has rank  $r = 1$ :

$$A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}.$$

- Find its largest singular value  $\sigma$  from  $A^T A$ .
- From its column space and row space, respectively, find unit vectors  $u$  and  $v$  so that

$$Av = \sigma u, \quad \text{and} \quad A = u\sigma v^T.$$

- From the nullspaces of  $A$  and  $A^T$  put numbers into the full SVD (Singular Value Decomposition) of  $A$ :

$$A = \begin{bmatrix} | & | \\ u & \dots \\ | & | \end{bmatrix} \begin{bmatrix} \sigma & 0 \\ 0 & \dots \end{bmatrix} \begin{bmatrix} | & | \\ v & \dots \\ | & | \end{bmatrix}^T.$$

*This page intentionally blank.*