

1 (a) $|A| = 2$ times the 3 by 3 determinant $= 2$ times $0 = 0$.

(b) A has rank 3 so we want three orthogonal basis vectors A, B, C :

$$\begin{aligned} A &= \text{first column } (1, 1, 1, 0) \\ B &= (\text{second column}) - (\text{projection onto first column}) \\ &= (-1, 1, 3, 0) - (1, 1, 1, 0) \cdot \frac{3}{3} \\ &= (-2, 0, 2, 0) \quad \text{check: orthogonal to first column} \\ C &= \text{last column } (0, 0, 0, 2) \end{aligned}$$

To orthogonalize divide by lengths:

$$\begin{aligned} q_1 &= \frac{A}{\|A\|} = \frac{A}{\sqrt{3}} \\ &= (1, 1, 1, 0)/\sqrt{3} \end{aligned}$$

$$\begin{aligned} q_2 &= \frac{B}{2\sqrt{2}} \\ &= (-1, 0, 1, 0)/\sqrt{2} \end{aligned}$$

$$\begin{aligned} q_3 &= \frac{C}{2} \\ &= (0, 0, 0, 1) \end{aligned}$$

(c) Adding 1 to the a_{11} entry will add its cofactor to the determinant:

$$\text{Cofactor } C_{11} = \begin{vmatrix} 1 & 5 & 0 \\ 3 & 9 & 0 \\ 0 & 0 & 2 \end{vmatrix} = -12.$$

2 (a)

$$\begin{bmatrix} 1 & -4 \\ 1 & -3 \\ 1 & -2 \\ 1 & -1 \\ 1 & 0 \\ 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} C \\ D \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 10 \end{bmatrix}$$

(b)

$$A^T A = \begin{bmatrix} 9 & 0 \\ 0 & 60 \end{bmatrix} \quad A^T b = \begin{bmatrix} 10 \\ 40 \end{bmatrix}$$

$$\text{Solve } A^T A \begin{bmatrix} C \\ D \end{bmatrix} = A^T b \text{ to find } C = \frac{10}{9}, D = \frac{40}{60}.$$

(c) The columns of A are a basis for the subspace. The projection is

$$p = C (\text{column 1}) + D (\text{column 2}).$$

3 (a) Q has rank n (the n orthonormal) columns are independent).

(b) $P = Q(Q^T Q)^{-1} Q^T = Q Q^T$.

(c) Check $P^T = P$: $(Q Q^T)^T = Q Q^T$.
Check $P^2 = P$: $(Q^T Q) Q^T = Q Q^T$.

4 (a) The length of λx is $|\lambda| \|x\|$.

The length squared of Qx is $(Qx)^T (Qx) = x^T Q^T Q x = x^T x = \|x\|^2$.

Thus $|\lambda| \|x\| = \|x\|$ and $|\lambda| = 1$.

Note: We did not use the correct notation when λ and x are complex. The reasoning stays the same.

(b) Projection onto the last column:

$$p = a \frac{a^T b}{a^T a} = \begin{bmatrix} 1 \\ -1 \\ -1 \\ 1 \end{bmatrix} 0 = \text{zero vector}.$$

Projection onto column space (which is all of R^4) is b itself.

(c) $|H_2 - \lambda I| = \begin{vmatrix} 1 - \lambda & 1 \\ 1 & -1 - \lambda \end{vmatrix} = \lambda^2 - 2 = 0$.

The eigenvalues are $\sqrt{2}$ and $-\sqrt{2}$. Check trace = 0 and determinant = -2.