

Your name is: _____

Grading 1

2

3

4

Please circle your recitation:

- | | | | | | | | |
|---------|-------|-------|--------------|---------|------|-------|------------|
| 1) Mon | 2-3 | 2-131 | S. Kleiman | 5) Tues | 12-1 | 2-131 | S. Kleiman |
| 2) Mon | 3-4 | 2-131 | S. Hollander | 6) Tues | 1-2 | 2-131 | S. Kleiman |
| 3) Tues | 11-12 | 2-132 | S. Howson | 7) Tues | 2-3 | 2-132 | S. Howson |
| 4) Tues | 12-1 | 2-132 | S. Howson | | | | |

1 (27 pts.) Suppose $A = \begin{bmatrix} .7 & .4 \\ .3 & .6 \end{bmatrix}$.

(a) Find the matrices Λ and S in the diagonalization formula $S^{-1}AS = \Lambda$.

(b) Find the matrix A^k (all four entries of the k^{th} power of A).

(c) Find the limit as $k \rightarrow \infty$ of $u_k = A^k u_0$ if $u_0 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

2 (25 pts.) Suppose a 3 by 3 real symmetric matrix A has eigenvalues 4, 4, 0.

(a) Find the determinant from the eigenvalues of $(2A - I)^{-1}$.

(b) True or false or not enough information:

This matrix A has 3 independent eigenvectors and can be diagonalized.

(c) True or false or not enough information:

The function $x^T Ax$ is never negative for any vector x .

(d) True or false or not enough information:

The matrix $\frac{1}{4}A$ is a Markov matrix.

(e) If A has orthonormal eigenvectors q_1, q_2, q_3 with $\lambda = 4, 4, 0$, find a formula for A in terms of q_1, q_2, q_3 using diagonalization.

3 (24 pts.) Suppose that A is an invertible 3 by 3 matrix.

- (a) Show me how to prove that $x^T A^T A x$ is *always positive* if x is not the zero vector. Why will this fail if A is not invertible?
- (b) Show me how to prove that $A^T A$ is *similar* to AA^T . Does it follow that these matrices have the same eigenvalues and eigenvectors?
- (c) If the SVD is written in the usual form $A = U\Sigma V^T$, what is the matrix $A^T A$ (reduced to the simplest form)?

- 4 (24 pts.)
- (a) Find the eigenvalues of $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ and $R = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$. Then find the eigenvalues of $A = \begin{bmatrix} a & 1 \\ 1 & a \end{bmatrix}$ and $R = \begin{bmatrix} b & -1 \\ 1 & b \end{bmatrix}$. The numbers a and b are real.
- (b) Under what condition on “ a ” do all solutions of $du/dt = Au$ approach zero as $t \rightarrow \infty$?
- (c) Under what conditions on “ b ” do all solutions of $dv/dt = Rv$ approach zero as $t \rightarrow \infty$?
- (d) Under what condition on “ a ” is the matrix A positive definite?